

PATIENT RECORD


SI joint pain diary

Referring physician _____

Date & time of appointment _____

Mark the figure with an "x" everywhere you have pain.

Referring Physician _____
Date & Time of Appointment _____
Please mark the figure with an "x" everywhere you have pain.



Time of Injection _____

Worst pain imaginable _____ SI joint pain _____

Do not use pencil, sharp object, or any other instrument to mark the figure.

Time	Before	30 min	1 hr	90 min	2 hr	3 hr	4 hr
10	○	○	○	○	○	○	○
9	○	○	○	○	○	○	○
8	○	○	○	○	○	○	○
7	○	○	○	○	○	○	○
6	○	○	○	○	○	○	○
5	○	○	○	○	○	○	○
4	○	○	○	○	○	○	○
3	○	○	○	○	○	○	○
2	○	○	○	○	○	○	○
1	○	○	○	○	○	○	○
No pain	○	○	○	○	○	○	○

Remarks _____

TIME OF
INJECTION

WORST EVER
SI JOINT PAIN

/10

SI JOINT PAIN
TODAY

/10

For each time period, shade the circle that corresponds to your SI joint pain level, then add any remarks below.

		AFTER INJECTION						
		Before	30 min	1 hr	90 min	2 hr	3 hr	4 hr
Worst pain imaginable	10	○	○	○	○	○	○	○
	9	○	○	○	○	○	○	○
	8	○	○	○	○	○	○	○
	7	○	○	○	○	○	○	○
	6	○	○	○	○	○	○	○
	5	○	○	○	○	○	○	○
	4	○	○	○	○	○	○	○
	3	○	○	○	○	○	○	○
	2	○	○	○	○	○	○	○
	1	○	○	○	○	○	○	○
No pain	0	○	○	○	○	○	○	○

Remarks _____

— NEXT STEPS

If sacroiliac joint pain limits your life, the SacroLox® procedure can help you *move with comfort again*.

Ask your healthcare provider if this minimally invasive option is right for you.

For more information, talk with your physician or visit your clinic's website.

Si-Logix | SacroLox
A Division of Evologics

www.evologicsamerica.com

This brochure is for informational purposes only and does not provide medical advice. Indications and contraindications: refer to the Instructions for Use (IFU). Individual results may vary. No outcome guarantees are expressed or implied.

Si-Logix | SacroLox
A Division of Evologics

SACROILIAC JOINT PAIN

Take back
your
movement

A minimally invasive approach to SI joint stabilization, designed to help you stand, sit and sleep more comfortably.



FOR PATIENTS


Understanding *SI joint pain*

01 Your sacroiliac joints

Connect the base of your spine (sacrum) to your pelvis. When these joints become irritated or unstable, it can cause pain that affects daily life.

02 What SI joint pain feels like

- Aching or sharp pain in the lower back, buttocks, or hips
- Discomfort that radiates to the groin or legs
- Pain that worsens with standing or walking
- Difficulty sitting for long periods or sleeping comfortably



Lateral view — the sacroiliac joint transfers load between the spine and pelvis.


Common experiences & *symptoms*

People with SI joint dysfunction can experience pain, weakness, and limited mobility.

COMMON SYMPTOMS

- Lower back pain (below L5)
- Pelvis or buttock pain
- Hip or groin pain
- Leg weakness or instability
- Numbness or tingling in the legs
- Pain when moving from sitting to standing
- Trouble sleeping due to pain
- Difficulty sitting for long periods

Your doctor can confirm SI joint dysfunction with a specific examination.



Posterior view — implant placement sites at the sacroiliac joints.

About the *SacroLox procedure*

PROCEDURE HIGHLIGHTS

- Minimally invasive approach
- Two cylindrical allograft implants inserted through a bone access needle
- Provides immediate joint stabilization and promotes fusion
- Usually performed under sedation in less than an hour

AFTER THE PROCEDURE

- Most patients walk the same day
- Return to light or non-strenuous work in about two weeks
- Follow your healthcare provider's postoperative instructions carefully



Two cylindrical allograft implants, shown beside a standard pencil for scale.